

Assessment of the variation in methods used by state agencies for collecting and processing benthic macroinvertebrate samples

James L. Carter¹ and Vincent H. Resh²

¹U.S. Geological Survey, 345 Middlefield Road, MS 465, Menlo Park, California 94025 USA

²University of California, Berkeley, Environmental Science, Policy, and Management, 201 Wellman Hall, Berkeley, California 94702 USA

Biographical Sketches of Authors

Jim Carter is an aquatic ecologist with the National Research Program, Water Resources Discipline of the U.S. Geological Survey. He studies the influence of physical and chemical factors on the composition and structure of benthic invertebrate assemblages in streams.

Vince Resh is a Professor of Entomology at the University of California, Berkeley, and has taught there for 25 y. He has done extensive research on stream and river bioassessment using macroinvertebrates.

Abstract

A survey of methods used by US state agencies for collecting and processing benthic macroinvertebrate samples from streams was conducted by questionnaire. The responses evaluated represent approximately 13,000-15,000 samples collected and processed per year. Kicknet devices are used in 64.5% of the methods. Mesh sizes vary among programs and within US EPA regions, but 80.2% use a mesh size between 500 and 600 mm. "Expert opinion" instead of random placement of the sampler is used by 70.6% of the methods, possibly making data obtained operator-specific. Only 26.3% of the methods sort all the organisms from a sample, the remainder subsample in the laboratory with most removing 100 organisms (range = 100-550). The magnification used for sorting ranges from 1× to 30×, which results in inconsistent separation of macroinvertebrates from detritus. Large/rare organisms are sorted by 53% of the methods, influencing estimates of richness. The taxonomic level used for identifying organisms varies among taxa; Ephemeroptera, Plecoptera, and Trichoptera are generally identified to a finer taxonomic resolution (genus and species) than other taxa. Although most programs use similar techniques, there currently exists a large range in how these techniques are applied, this would make calibration among programs challenging. Limited testing could be designed to evaluate whether these differences affect data comparability and, more importantly, determining levels of environmental impairment. A companion survey to evaluate methods used for data analysis is currently being finalized.